

17-21 September 2018,  
Le Corum, Montpellier - France

# Eucalyptus 2018



Managing  
*Eucalyptus* plantations  
under global changes



Abstracts Book

# Using multispecies NIRS calibration for predicting chemical properties of eucalypts wood

Andriambelo Radonirina Razafimahatratra <sup>\*† 1</sup>, Tahiana Ramananantoandro <sup>1</sup>, Sophie Nourissier <sup>2</sup>, Zo Elia Mevanarivo <sup>1</sup>, Mario Tomazello Filho <sup>3</sup>, Garel Makouanzi <sup>4</sup>, Anne Clément-Vidal <sup>2</sup>, José Rodrigues <sup>5</sup>, Chaix Gilles <sup>2</sup>

<sup>1</sup> Ecole Supérieure des Sciences Agronomiques - Mention Foresterie et Environnement (ESSA-Forets) – BP 175, Antananarivo 101, Madagascar

<sup>2</sup> CIRAD (CIRAD) – AGAP – Cirad Campus of Lavalette TA A-108/01 34398 Montpellier Cedex 5, France

<sup>3</sup> Escola Superior de Agricultura "Luiz de Queiroz" (ESALQ) – scola Superior de Agricultura "Luiz de Queiroz" Avenida Pádua Dias, 11 - Piracicaba/SP - CEP 13418-900, Brazil

<sup>4</sup> Ecole Supérieure des Sciences Agronomiques et Forestières (ENSAF) – Université Marien NGOUABI, Congo - Brazzaville

<sup>5</sup> Instituto Superior de Agronomia (ISA) – University of Lisboa, Portugal

Breeding programs in Africa are generally based on growth criteria and rarely on wood chemical properties. Indeed, chemical analysis are often expensive, time-consuming and require several replicates. Then, using NIRS to predict these properties is a relevant solution. The research question focuses on the possibility of using multispecies models to predict properties of different species. This study considers 7 chemical properties (extractives, Klason lignin, acid-soluble lignin ASL, SG ratio, holocellulose, alpha-cellulose, hemicelluloses) based on 367 samples from 4 countries, belonging to 5 eucalypt species with hybrids (*E. robusta*, *camaldulensis*, *urophylla*, *uropellita*, *urograndis*). Established models were validated by cross- and test-set validation. Results shows that all R<sup>2</sup>CV are greater than 0.73, and all %RMSECV are less than 8.3% except for extractives and ASL. Prediction errors (%RMSEP) are always less than 9.5% except for these 2 properties, with respectively 23.6% and 18.1%. Prediction errors are always less than the double of the error of laboratory (%SEL). This study shows that multispecies NIRS models can be used to predict chemical properties, there is no significant difference between measurement error obtained with standardized method and %RMSEP. This method is particularly well-suited for a rapid wood phenotyping of multiple samples belonging to different species.

**Keywords:** Near InfraRed Spectroscopy, multispecies prediction model, error of laboratory, chemical properties, Eucalyptus

---

\*Speaker

†Corresponding author: andriambelo.radonirina@gmail.com